

Origin of G-type Antiferromagnetism and Orbital-Spin Structures in LaTiO_3

Masahito MOCHIZUKI and Masatoshi IMADA

*Institute for Solid State Physics, University of Tokyo,
5-1-5 Kashiwa-no-ha, Kashiwa, Chiba 277-8581*

(Received February 1, 2008)

The possibility of the D_{3d} distortion of TiO_6 octahedra is examined theoretically in order to understand the origin of the G-type antiferromagnetism (AFM(G)) and experimentally observed puzzling properties of LaTiO_3 . By utilizing an effective spin and pseudospin Hamiltonian with the strong Coulomb repulsion, it is shown that AFM(G) state is stabilized through the lift of the t_{2g} -orbital degeneracy accompanied by a tiny D_{3d} -distortion. The estimated spin-exchange interaction is in agreement with that obtained by the neutron scattering. Moreover, the level-splitting energy due to the distortion can be considerably larger than the spin-orbit interaction even when the distortion becomes smaller than the detectable limit under the available experimental resolution. This suggests that the orbital momentum is fully quenched and the relativistic spin-orbit interaction is not effective in this system, in agreement with recent neutron-scattering experiment.

KEYWORDS: LaTiO_3 , GdFeO_3 -type distortion, D_{3d} -crystal field, orbital ordering, second-order perturbation theory, G-type antiferromagnetism

§1. Introduction

Perovskite-type Ti oxide $RTiO_3$ have recently attracted much interest because of the rich magnetic and orbital phases caused by an interplay of spin and orbital degrees of freedom.¹⁾ In these compounds, Ti^{3+} has a $3d^1$ configuration, and one of the threefold t_{2g} -orbitals is occupied at each transition-metal site. The crystal structure is a pseudocubic perovskite with GdFeO_3 -type distortion in which the TiO_6 octahedra forming the perovskite lattice tilt alternately. The unit cell contains four TiO_6 -octahedra, as shown in Fig. 1. With a smaller ionic radius of the R ion, the distortion increases with the decrease in the Ti-O-Ti




Fig. 1. GdFeO₃-type distortion.

bond angle from 180°. The bond angle can be controlled by use of the solid-solution systems $\text{La}_{1-y}\text{Y}_y\text{TiO}_3$ or in $R\text{TiO}_3$, by varying the R ions.

Recently, the magnetic phase diagrams have been investigated intensively in the plane of temperature and the magnitude of the distortion.²⁻⁵⁾ In particular, LaTiO_3 exhibits a G-type AFM (AFM(G)) ground state with magnetic moment of $0.45 \mu_B$,⁴⁾ and the Néel temperature (T_N) takes about 130 K. With increasing the GdFeO₃-type distortion, T_N decreases rapidly and is suppressed to almost zero, subsequently a ferromagnetic (FM) ordering appears. In Y-rich systems and in YTiO_3 in which the GdFeO₃-type distortion is relatively large, the system shows a FM ground state accompanied by a large Jahn-Teller (JT) distortion. Recently, the AFM-FM phase transition with the second-order-like behavior has been understood from the strong two-dimensionality in the spin-coupling realized near the transition point.^{6,7)} However, the origin and nature of AFM(G) state in the end compounds LaTiO_3 remains unresolved.

Recent model Hartree-Fock study predicts the AFM(G) state with the spin-orbit ground state in the absence of static JT distortion.^{8,9)} However, in contrast to the prediction from the spin-orbit ground state, recent neutron-scattering study shows the spin-wave spectrum well described by an isotropic spin-1/2 Heisenberg model with a nearest-neighbor superexchange constant $J \sim 15.5 \text{ meV}$.¹¹⁾ This also suggests the absence of unquenched orbital moments. When the orbital degeneracy remains, a FM spin structure with antiferro-orbital ordering in which the neighboring orbitals are orthogonal to each other is eventually expected in the presence of the transfers and the Coulomb-exchange interaction. Indeed, recent weak coupling study shows that in the cubic-crystal field in the absence of the static JT distortion, a FM state, out of which two states $\frac{1}{\sqrt{2}}(yz + izx) \uparrow$ and $xy \uparrow$ are alternating is favored both by the spin-orbit interaction and by the spin-orbital superexchange interaction.¹⁰⁾ However, no evidence for the orbital ordering is found in a resonant x-ray scattering study.¹¹⁾

Recently, a possible orbital liquid state was proposed on the basis of small exchange interaction in the orbital sector in the AFM(G) state.¹²⁾ However, in a system with strong orbital




Fig. 2. D_{3d} distortion.

fluctuations, the spin and orbital degrees of freedom strongly couple, and both degrees of freedom can not be determined independently to each other. In this circumstance, the origin of AFM(G) state in LaTiO_3 is to be clarified in a self-consistent manner.

In this letter, in order to understand the above properties of AFM(G) state in LaTiO_3 , we propose that the puzzle is solved if we assume the existence of a small D_{3d} -crystal field. If the D_{3d} distortion exists, TiO_6 -octahedron is contracted along the threefold direction as shown in Fig. 2. As a result, threefold degenerate t_{2g} -levels split into non-degenerate lower a_{1g} -level (φ_1) and twofold-degenerate higher e_g -levels (φ_2, φ_3). In the distortion with $[1,1,1]$ -trigonal axis, the a_{1g} representation is $\frac{1}{\sqrt{3}}(xy + yz + zx)$ and the e_g representations are $\frac{1}{\sqrt{6}}(2xy - yz - zx)$ and $\frac{1}{\sqrt{2}}(yz - zx)$. Since the lowest level has no degeneracy, this distortion significantly and sensitively lowers the energy of the electron system in t_{2g}^1 configuration. Though the d -type JT distortion is stabilized in YTiO_3 or in the compounds with large GdFeO_3 -type distortion due to the effects of covalency between O $2p$ and unoccupied d -orbitals on the R -site cations,¹³⁾ LaTiO_3 with small GdFeO_3 -type distortion turns out to be unstable to the D_{3d} distortion. The magnitude of the distortion can be denoted by the D_{3d} -distortion angle ω (see Fig. 2). By utilizing the effective spin and pseudospin Hamiltonian in the strong Coulomb repulsion, we solve the self-consistent Hartree-Fock equation and show that the emergence of the AFM(G) state and the following features in LaTiO_3 are well explained by the effects of D_{3d} distortion even when ω is as small as 1° :

- $T_N \sim 130$ K, $J \sim 15.5$ meV,
- magnetic moment of $0.45 \mu_B$,
- no detectable JT distortion,
- no evidence for orbital ordering in the resonant x-ray scattering.

We analyze why the orbital ordering has not been detected so far, and propose experimental ways for testing the validity of our proposal.

§2. Formalism

We start with the multiband d - p model in which the full degeneracies of Ti 3*d* and O 2*p* orbitals as well as the on-site Coulomb and exchange interactions are taken into account.⁷⁾ The nearest-neighbor d - p and p - p transfers are given in terms of Slater-Koster parameters $V_{pd\pi}$, $V_{pd\sigma}$, $V_{pp\pi}$ and $V_{pp\sigma}$.¹⁴⁾ The on-site Coulomb interactions are expressed using Kanamori parameters, u , u' , j and j' which satisfy the following relations:^{15, 16)} $u = U + \frac{20}{9}j$, $u' = u - 2j$ and $j = j'$. Here, U gives the magnitude of the multiplet-averaged d - d Coulomb interaction. The charge-transfer energy Δ , which describes the energy difference between occupied O 2*p* and unoccupied Ti 3*d* levels, is defined by using U and energies of the bare Ti 3*d* and O 2*p* orbitals ε_d^0 and ε_p as $\Delta = \varepsilon_d^0 + U - \varepsilon_p$, since the characteristic unoccupied 3*d* level energy on the singly occupied Ti site is $\varepsilon_d^0 + U$. The values of the parameters are estimated by the cluster-model analyses of valence-band and transition-metal 2*p* core-level photoemission spectra and the analyses of the first-principle band calculations.^{17, 18)} We take the values of these parameters as $\Delta = 5.5$ eV, $U = 4.0$ eV, $V_{pd\sigma} = -2.4$ eV, $V_{pd\pi} = 1.3$ eV, $V_{pp\sigma} = 0.52$ eV, $V_{pp\pi} = -0.11$ eV and $j = 0.46$ eV throughout the present calculation. The effects of the GdFeO₃-type distortion are considered through the transfer integrals which is defined by using the Slater-Koster's parameters¹⁴⁾. In the present calculations, we simulate the GdFeO₃-type distortion by rotating the TiO₆ octahedra by angle $+14.1^\circ$ and -14.1° around the $[1,1,1]$ and $[-1,-1,1]$ axes. As a result, the Ti-O-Ti bond angle takes $\sim 157^\circ$. The effects of the D_{3d} distortion are also considered. We integrate out the O 2*p* orbital degrees of freedom in the path-integral formalism to arrive at the effective multiband Hubbard model which includes only Ti 3*d* orbital degrees of freedom. Starting with thus obtained multiband Hubbard Hamiltonian, we can derive an effective Hamiltonian in the low-energy region on the subspace of states with singly occupied t_{2g} orbitals at each transition-metal site by utilizing a second-order perturbation theory. The states of 3*d* electrons localized at the transition-metal sites can be represented by two quantum numbers, the z -component of the spin S_z and the type of the occupied orbitals. The threefold degeneracy of t_{2g} -orbitals at each site can be described by a pseudospin with the spin-1 operator $\boldsymbol{\tau}$. We follow an approach similar to the Kugel-Khomskii formulation.^{20, 21)} We express the 3*d* electron operators in terms of $\boldsymbol{S}$ and $\boldsymbol{\tau}$ to arrive at the effective spin and pseudospin Hamiltonian: $H_{\text{eff}} = H_{\text{cry.}} + H_{t_{2g}} + H_{e_g}$. The first term $H_{\text{cry.}}$ denotes the effects of D_{3d} -crystal field. This term is obtained from the zeroth-order perturbational processes. The second term $H_{t_{2g}}$ is obtained from the second-order perturbational processes whose intermediate states contain only t_{2g} -orbital degrees of freedom. The third term H_{e_g} is obtained from the second-order perturbational processes whose intermediate states contain e_g -orbital degrees of freedom. We neglect the relativistic spin-orbit coupling in the model and discuss this issue later.




Fig. 3. Energies of each magnetic structures with config. 1 as functions of the D_{3d} -distortion angle. See the text for details.

§3. Results

There are four trigonal-axes in the TiO_6 octahedron, namely, $[1,1,1]$, $[1,1,-1]$, $[1,-1,1]$ and $[1,-1,-1]$ with respect to the x , y and z axes. At this stage, there are several possibilities for configurations of the trigonal axes at sites 1, 2, 3, and 4. The spin-couplings between the neighboring sites depend on the configuration. There are five configurations which lower the AFM(G) spin structure in energy as shown in Table I. In this letter, we examine the AFM(G) state with a trigonal-axes configuration in which $[1,1,1]$ -axis is chosen at every sites (config. 1), and we show that a considerably small D_{3d} -distortion does strongly stabilize the AFM(G) state and the experimentally observed properties are well explained when we assume the distortion. It is an interesting problem to study which configuration has the lowest energy, which would require an entirely separate work and is left for future discussion. In the D_{3d} distortion with config. 1, the lowest level is $\frac{1}{\sqrt{3}}(xy + yz + zx)$ at every sites. In Fig. 3, we plot the energies of several magnetic structures with config.1 as functions of D_{3d} -distortion angle ω . In the region of $\omega > 1^\circ$, the AFM(G) state is strongly stabilized relative to other structures. This may indeed drive the D_{3d} distortion even in the cost of lattice energy. Since the lattice elastic-energy is not accounted in Fig. 3, the real energy minimum including the elastic-energy cost should be in the range $0^\circ < \omega < 8^\circ$. In Fig. 4, we plot the t_{2g} -level-splitting energy due to the D_{3d} -crystal field ($\Delta_{D_{3d}}$) as a function of ω . In the region of $\omega > 1^\circ$, the value of $\Delta_{D_{3d}}$ is sufficiently larger than the coupling constant of the relativistic spin-orbit interaction in Ti^{3+} ion ($\xi_d = 0.018$ eV),²²⁾ though this distortion angle $\omega > 1^\circ$ is substantially smaller than the tilting angle of GdFeO_3 -type distortion. In fact, it is not surprising if such small D_{3d} distortion has not been detected so far.

When the level-splitting energy is sufficiently large as compared to the characteristic energy of the spin-orbital exchange interaction, the orbital occupation at each site is restricted to

Table I. Trigonal-axes configurations which lower the AFM(G) spin structure in energy.

	site 1	site 2	site 3	site 4
config.1	[1,1,1]	[1,1,1]	[1,1,1]	[1,1,1]
φ 1	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$
config.2	[1,-1,1]	[1,-1,1]	[1,-1,1]	[1,-1,1]
φ 1	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$
config.3	[1,1,1]	[1,1,1]	[1,1,-1]	[1,1,-1]
φ 1	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy - yz - zx)$	$\frac{1}{\sqrt{3}}(xy - yz - zx)$
config.4	[1,1,-1]	[1,1,-1]	[1,1,1]	[1,1,1]
φ 1	$\frac{1}{\sqrt{3}}(xy - yz - zx)$	$\frac{1}{\sqrt{3}}(xy - yz - zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$	$\frac{1}{\sqrt{3}}(xy + yz + zx)$
config.5	[1,-1,1]	[1,-1,1]	[1,-1,-1]	[1,-1,-1]
φ 1	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$	$\frac{1}{\sqrt{3}}(-xy - yz + zx)$	$\frac{1}{\sqrt{3}}(-xy + yz + zx)$	$\frac{1}{\sqrt{3}}(-xy + yz + zx)$




Fig. 4. Level-splitting energy due to the D_{3d} -crystal field ($\Delta_{D_{3d}}$) as a function of the D_{3d} -distortion angle.




Fig. 5. Characteristic transfers which contribute to the spin-exchange interaction.




Fig. 6. Spin-exchange interactions estimated under the assumption that orbital occupation is restricted to the lowest level (J_{Heis}^{ab} and J_{Heis}^c) are plotted. Inset shows those calculated using other parameters within the error bars of the analyses of photoemission spectra. See the text for details.

$\frac{1}{\sqrt{3}}(xy + yz + zx)$ independently of the magnetic structure. At this stage, we can estimate the value of the spin-coupling constant in the ab -plane (J_{Heis}^{ab}) and that along the c -axis (J_{Heis}^c) under this assumption. In Fig. 5, the characteristic transfers which contribute to the spin-exchange interaction are presented. Using the notation of transfer integrals as shown in Fig. 5, J_{Heis}^{ab} and J_{Heis}^c are expressed in the strong coupling expansion as,

$$J_{\text{Heis}}^{ab}, J_{\text{Heis}}^c = 4 \frac{t_1^2}{u} - 2 \frac{(t_2^2 + t_3^2)j}{(u' + \Delta_{D3d})^2} - 2 \frac{(t_2'^2 + t_3'^2)j}{(u' + \Delta_{D3d})^2}. \quad (1)$$

In Fig. 6, the values of J_{Heis}^{ab} and J_{Heis}^c are plotted as functions of ω . Both J_{Heis}^{ab} and J_{Heis}^c take approximately 0.015-0.022 eV, and they are in agreement with the value obtained by the neutron scattering experiment of ~ 15.5 meV. In LaTiO_3 , the magnetic moment takes $0.45 \mu_B$, which is reduced from the spin-only moment. Recent optical measurement shows that LaTiO_3 has a considerably small optical gap of ~ 0.1 eV in the vicinity of the metal-insulator phase boundary with strong itinerant character.²⁾ Therefore, in this system, we expect that

some amount of orbital and spin fluctuations remain. The reduction of the magnetic moment may easily be attributed to these fluctuations. We note that in this context, the calculated J_{Heis}^{ab} and J_{Heis}^c in the localized limit seem rather small at first sight. To elucidate this issue, we have examined how the values of J_{Heis}^{ab} and J_{Heis}^c change within the error bars of the parameters estimated from the analyses of photoemission spectra. In LaTiO_3 with strong itinerant character, we can expect that the Coulomb and exchange interactions are rather small. We calculate the values in the case of $U = 3.0$ eV, $j = 0.50$ eV and $\Delta = 5.0$ eV and in the case of $U = 3.0$ eV, $j = 0.40$ eV and $\Delta = 5.0$ eV. The values take ~ 0.03 eV, which are larger than the previous estimate (see Inset of Fig. 6). Considering these variations and uncertainties, we conclude that the calculated J_{Heis}^{ab} and J_{Heis}^c in the localized limit are in qualitative agreement with the experimentally obtained J in the presence of itinerant charge fluctuations.

Here, a question arises: why has the orbital ordering caused by the occupation of the lowest level in the D_{3d} distortion not been detected experimentally so far? Detection of the orbital ordering by the resonant x-ray scattering is based on the splitting of the Ti $4p$ levels induced by the orbital ordering of the Ti $3d$ states. When the Ti $3d$ orbitals are ordered in the D_{3d} distortion, the Coulomb interaction between Ti $4p$ and occupied $3d$ orbitals, U_{4p-3d} and the interaction between Ti $4p$ and ligand O $2p$ orbitals, U_{4p-2p} make opposite contributions. Among the threefold Ti $4p$ orbitals, the orbital directed along the trigonal axis has higher energy due to U_{4p-3d} . On the contrary, along this axis, U_{4p-2p} is weakened so that the crystal field works to lower the Ti $4p$ orbital. In addition, the weak hybridizations between the Ti $4p$ and O $2p$ orbitals might lower the Ti $4p$ orbital. This cancellation may lead to the absence of detectable orbital ordering in the resonant x-ray scattering in LaTiO_3 .¹¹⁾ The orbital ordering accompanied by the D_{3d} distortion would be more easily detected in the AFM(G) compounds with relatively large optical gap and small charge fluctuations: PrTiO_3 and NdTiO_3 .

In summary, in order to interpret the experimentally observed properties of AFM(G) state in LaTiO_3 , we have examined the effects of possible D_{3d} -crystal field as a candidate for their origin. In the presence of the D_{3d} -crystal field, threefold cubic- t_{2g} -levels are split into non-degenerate a_{1g} - and twofold-degenerate e_g -levels. Since the lowest level has no degeneracy, this distortion strongly lowers the energy of the electron system in t_{2g}^1 configuration. Moreover, the calculated D_{3d} -splitting energy $\Delta_{D_{3d}}$ suggests that the D_{3d} -splitting dominates over the relativistic spin-orbit interaction in the region of D_{3d} -distortion angle $\omega > 1^\circ$. The calculated spin-exchange constant is in agreement with that obtained experimentally. Finally, our proposal for the small D_{3d} distortion should in principle be observed by the displacement of O ions if the experimental resolution is sufficient.

M. M. thanks H. Asakawa and T. Mizokawa for useful comments. This work is supported by “Research for the Future Program” (JSPS-RFTF97P01103) from the Japan Society for the Promotion of Science.

-
- 1) For a review see M. Imada, A. Fujimori and Y. Tokura: Rev. Mod. Phys. **70** (1998) 1039.
 - 2) Y. Okimoto, T. Katsufuji, Y. Okada, T. Arima and Y. Tokura: Phys. Rev. B **51** (1995) 9581.
 - 3) T. Katsufuji, Y. Taguchi and Y. Tokura: Phys. Rev. B **56** (1997) 10145.
 - 4) J. P. Goral, J. E. Greedan and D. A. Maclean: J. Solid State Chem. **43** (1982) 244.
 - 5) J. E. Greedan: J. Less-Common Met. **111** (1985) 335.
 - 6) M. Mochizuki and M. Imada: J. Phys. Soc. Jpn. **69** (2000) 1982.
 - 7) M. Mochizuki and M. Imada: cond-mat/0102223 (to appear in J. Phys. Soc. Jpn.)
 - 8) T. Mizokawa and A. Fujimori: Phys. Rev. B **51** (1995) 12880.
 - 9) T. Mizokawa and A. Fujimori: Phys. Rev. B **54** (1996) 5368.
 - 10) M. Mochizuki and M. Imada: unpublished
 - 11) B. Keimer, D. Casa, A. Ivanov, J. W. Lynn, M. v. Zimmermann, J. P. Hill, D. Gibbs, Y. Taguchi and Y. Tokura: Phys. Rev. Lett. **85** (2000) 3946.
 - 12) G. Khaliullin and S. Maekawa: Phys. Rev. Lett. **54** (2000) 3950.
 - 13) T. Mizokawa, D. I. Khomskii and G. A. Sawatzky: Phys. Rev. B **60** (1999) 7309.
 - 14) J. C. Slater and G. F. Koster: Phys. Rev. **94** (1954) 1498.
 - 15) B. H. Brandow: Adv. Phys. **26** (1977) 651.
 - 16) J. Kanamori: Prog. Theor. Phys. **30** (1963) 275.
 - 17) T. Saitoh, A. E. Bocquet, T. Mizokawa and A. Fujimori: Phys. Rev. B **52** (1995) 7934.
 - 18) P. Mahadevan, N. Shanthi and D. D. Sarma: Phys. Rev. B **54** (1996) 11199.
 - 19) W. A. Harrison: Electronic Structure and the Properties of Solids (Dover, New York, 1989).
 - 20) K. I. Kugel and D. I. Khomskii: Pisma. Zh. Eksp. Teor. Fiz. **15** (1972) 629.
 - 21) K. I. Kugel and D. I. Khomskii: Zh. Eksp. Teor. Fiz. **64** (1973) 1429.
 - 22) S. Sugano, Y. Tanabe and H. Kamimura: Multiplets of Transition-Metal Ions in Crystals (Academic, New York 1970)